

Tehtävät palautetaan viimeistään keskiviikkona 19.11. Physicum 2. kerroksen A-siiven aulassa olevaan laatikkoon. HUOM! Maanantaina 10.11. ei pidetä harjoituksia, eikä keskiviikkona 12.11. luentoja! Ylimääräinen harjoitus ma 15.12. klo 10.

1. Erään kaksiulotteisen moniston erään koordinaattiympäristön kuva on  $\mathbf{R}^2$ . Anna käyrä, jonka tangenttivektori pisteessä jonka koordinaatit ovat  $(1,1)$  on  $X = a \frac{\partial}{\partial x} + b \frac{\partial}{\partial y}$ ,  $a, b$  reaalityyppiset luvut. Laske  $Xf$ , kun  $f$  on funktio  $f(x, y) = \tanh(3x + y^2)$ . Olkoot  $(\xi, \eta)$  koordinaatit toisessa koordinaattiympäristössä, johon piste  $(x, y) = (1, 1)$  myös kuuluu, ja olkoot transitiofunktiot  $\xi = \frac{x+1}{x+y}, \eta = \frac{y+1}{x+y}$  eräässä pisteen  $(1,1)$  ympäristössä. Laske valitsemasi käyrän kaava uusissa koordinaateissa. Muunna  $X$  koordinaattikantaan  $\frac{\partial}{\partial \xi}, \frac{\partial}{\partial \eta}$ , ja tarkista että se on käyräsi tangentti.

2. Johda muunnoskaava tensorin

$$T = T_{\nu_1 \nu_2 \nu_3}^{\mu_1 \mu_2} \frac{\partial}{\partial x^{\mu_1}} \otimes \frac{\partial}{\partial x^{\mu_2}} \otimes dx^{\nu_1} \otimes dx^{\nu_2} \otimes dx^{\nu_3}$$

komponenteille koordinaattimuunnoksessa  $x \rightarrow y$ .

3. Olkoon  $X \in \mathbf{R}^3$ :n vektorikenttä

$$X = X^i(\vec{x}) \frac{\partial}{\partial x^i}.$$

Määritellään  $(1,1)$ -tensorikenttä  $S_{st}$ ,

$$S_{st} = S_j^i \frac{\partial}{\partial x^i} \otimes dx^j,$$

jonka komponentit  $S_j^i$  ovat

$$S_j^i = \frac{1}{2} \frac{\partial X^i}{\partial x^j} + \frac{1}{2} \frac{\partial X^j}{\partial x^i} - \frac{1}{3} \left( \sum_{k=1}^3 \frac{\partial X^k}{\partial x^k} \right) \delta_j^i.$$

Laske komponentit  $S_j^i$  kun  $X^1 = 0, X^2 = k \cdot x^1, X^3 = 0$ ,  $k$  on vakio (reaalityyppinen). Komponentti:  $S_{st}$  on elastiseen kappaleeseen kohdistuvaa halkaisevaa jännitystä (shearing strain) kuvaava jännitystensori. Tehtävässä kappale on elastinen palkki, johon kohdistuu pituus-suunnassa halkaiseva jännitys. Katso K. Symon, *Mechanics*, s. 237, 437-8.

4. Olkoon  $f : M \rightarrow N$  ja  $g : N \rightarrow P$ . Osoita että yhdistetyn kuvauksen  $gf : M \rightarrow P$  työntökuvaus on  $(gf)_* = g_* f_*$ , mutta että sen takaisinvektori on  $(gf)^* = f^* g^*$ .

5. Lue kappale 4.4.8., Vektorikentän generoima virtaus, ja ratkaise seuraava tehtävä:

**Hamiltonin liikeyhtälöt esimerkkinä vektorikentän generoimasta virtauksesta.** Klassisesta mekaniikasta on (toivottavasti) tuttua se, että systeemiä jolla on  $N$  vapausastetta voidaan kuvata  $N$ :llä yleistetyllä koordinaatilla  $q_i$  ja kanonisella impulssilla  $p_i$ , missä  $i = 1, \dots, N$ . Yleistetyt koordinaatit ja kanoniset impulssit voidaan ajatella  $2N$ -ulotteisen moniston  $M$  koordinaateiksi,  $M$ :ää kutsutaan *faasiavaruudeksi*. Systemin

dynamiikkaa kuvaa Hamiltonin funktio  $H : M \rightarrow \mathbb{R}$ ,  $H = H(q_1, \dots, q_N, p_1, \dots, p_N)$  ja sen liikeyhtälöt voidaan kirjoittaa ensimmäisen asteen differentiaaliyhtälöryhmänä

$$\frac{dq_i}{dt} = \frac{\partial H}{\partial p_i}; \quad \frac{dp_i}{dt} = -\frac{\partial H}{\partial q_i}.$$

Em. yhtälöitä kutsutaan Hamiltonin yhtälöiksi. Kirjoitetaan seuraavaksi yhtälöt eritavalla. Määritellään vektorikenttä  $X_H$  faasiavaruudessa  $M$ ,

$$X_H = \sum_{i=1}^N \left\{ \frac{\partial H}{\partial p_i} \frac{\partial}{\partial q_i} - \frac{\partial H}{\partial q_i} \frac{\partial}{\partial p_i} \right\}.$$

Vektorikentän  $X_H$  integraalikäyrä  $x_H(t) = (q_1(t), \dots, q_N(t), p_1(t), \dots, p_N(t))$  on käyrä monistolla  $M$ .

- i) Osoita että integraalikäyriä  $x_H(t)$  kuvaavat yhtälöt vastaavat Hamiltonin yhtälöitä.
- ii) Olkoon  $M = \mathbb{R}^2$ , eli  $N = 1$ , ja  $H = \frac{1}{2}(p^2 + q^2)$ . Etsi  $X_H$  ja sen generoima virtaus  $\sigma(t, x_0)$  missä  $x_0 = (q_0, p_0) = (1, 0)$ . Piirrä kuva.
- iii) Samoin kuin edellä mutta nyt  $H = \frac{1}{2}(p^2 - q^2)$  ja  $x_0 = (1, 1)$ .
- iv) Nyt  $M = T^2$ , koordinaatteina  $q, p \in [0, 2\pi]$ , ja  $H = \cos(p)$ . Etsi integraalikäyrän yhtälö. Piirrä kuva integraalikäyrästä  $T^2$ :lla.

English text:

Return your solutions to the box in the entrance hall on the second floor of the A-wing in Physicum by Wednesday, November 19. NOTE: No examples class on Monday, November 10 and no lectures on Wednesday, November 12!

1. The image of a coordinate patch of a certain two-dimensional manifold is  $\mathbf{R}^2$ . Give one curve, whose tangent vector in the point with coordinates  $(1,1)$  is  $X = a \frac{\partial}{\partial x} + b \frac{\partial}{\partial y}$ ,  $a, b$  real. Give the value of  $X$  on the function given by  $f(x, y) = \tanh(3x + y^2)$ . Let  $(\xi, \eta)$  be the coordinates on another patch, to which  $(x, y) = (1, 1)$  also belongs, and let the transition functions be  $\xi = \frac{x+1}{x+y}$ ,  $\eta = \frac{y+1}{x+y}$  in a neighbourhood around  $(1,1)$ . Compute the form of the curve you chose in these new coordinates. Transform  $X$  into the coordinate basis  $\frac{\partial}{\partial \xi}, \frac{\partial}{\partial \eta}$ , and check that it is the tangent vector of your curve.

2. Derive the transformation rule for the components of the tensor

$$T = T^{\mu_1 \mu_2}_{\nu_1 \nu_2 \nu_3} \frac{\partial}{\partial x^{\mu_1}} \otimes \frac{\partial}{\partial x^{\mu_2}} \otimes dx^{\nu_1} \otimes dx^{\nu_2} \otimes dx^{\nu_3}$$

in the coordinate transformation  $x \rightarrow y$ .

3. Let  $X$  be a vector field in  $\mathbb{R}^3$ ,

$$X = X^i(\vec{x}) \frac{\partial}{\partial x^i}.$$

We then define a (1,1) tensor field  $S_{st}$ ,

$$S_{st} = S_j^i \frac{\partial}{\partial x^i} \otimes dx^j ,$$

with the components  $S_j^i$  given by

$$S_j^i = \frac{1}{2} \frac{\partial X^i}{\partial x^j} + \frac{1}{2} \frac{\partial X^j}{\partial x^i} - \frac{1}{3} \left( \sum_{k=1}^3 \frac{\partial X^k}{\partial x^k} \right) \delta_j^i .$$

Calculate the components  $S_j^i$  when  $X^1 = 0, X^2 = k \cdot x^1, X^3 = 0$ ,  $k$  is a (real valued) constant. Comment:  $S_{st}$  is a shear tensor, which describes the shearing strain on an elastic body. In this exercise the body is an elastic beam, subject to a shearing force parallel to the beam. See K. Symon, *Mechanics*, s. 237, 437-8.

4. Let  $f : M \rightarrow N$  and  $g : N \rightarrow P$ . Show that the differential map of the composite map  $gf : M \rightarrow P$  is  $(gf)_* = g_* f_*$ , but that the pullback of  $gf$  is  $(gf)^* = f^* g^*$ .

5. Read section 4.4.8, Flow Generated by a Vector Field, and solve the following problem:

**Hamilton's Equations as an Example of a Flow Generated by Vector Field.**

It should be familiar from Classical Mechanics that a system with  $N$  degrees of freedom can be described by generalized coordinates  $q_i$  and canonical momenta  $p_i$ , where  $i = 1, \dots, N$ . The generalized coordinates and canonical momenta can be thought as coordinates of a  $2N$ -dimensional manifold  $M$ , called the *phase space*. The dynamics of the system is given by the Hamiltonian,  $H : M \rightarrow R$ ,  $H = H(q_1, \dots, q_N, p_1, \dots, p_N)$ ; its equations of motion can be written as a group of first order differential equations

$$\frac{dq_i}{dt} = \frac{\partial H}{\partial p_i} ; \quad \frac{dp_i}{dt} = - \frac{\partial H}{\partial q_i} .$$

These are called the Hamilton's equations. We will next reformulate them in a different way. Let's define a vector field  $X_H$  in the phase space  $M$ ,

$$X_H = \sum_{i=1}^N \left\{ \frac{\partial H}{\partial p_i} \frac{\partial}{\partial q_i} - \frac{\partial H}{\partial q_i} \frac{\partial}{\partial p_i} \right\} .$$

The vector field  $X_H$  gives rise to integral curves  $x_H(t) = (q_1(t), \dots, q_N(t), p_1(t), \dots, p_N(t))$  on the manifold  $M$ .

i) Show that the equation defining the integral curves  $x_H(t)$  is equivalent to Hamilton's equations.

ii) Let  $M = R^2$ , i.e.  $N = 1$ , and  $H = \frac{1}{2}(p^2 + q^2)$ . Find  $X_H$  and the generated flow  $\sigma(t, x_0)$  where  $x_0 = (q_0, p_0) = (1, 0)$ . Illustrate it by a figure.

iii) As before, but now with  $H = \frac{1}{2}(p^2 - q^2)$  ja  $x_0 = (1, 1)$ .

iv) Now  $M = T^2$ , with coordinates  $q, p \in [0, 2\pi]$ , and  $H = \cos(p)$ . Find the equation of the integral curve. Draw a figure of the curves on  $T^2$ .